

Total No. of printed pages = 10

3 (Sem 1) MAT M1

2015

MATHEMATICS

(Major)

Paper : 1.1

Full Marks - 80

Time - Three hours

The figures in the margin indicate full marks for the questions.

1. Answer/choose the correct option: $1 \times 10 = 10$
- (a) Give an example of a relation on the set of real numbers \mathbb{R} which is reflexive and transitive but not symmetric.
- (b) Find all partitions of the set $x = \{1, 2, 3\}$.
- (c) Let Q_+ be the set of all positive rational numbers and $*$ be a binary operation on Q_+ defined by $a * b = \frac{ab}{3}$, $\forall a, b \in Q_+$. Find the identity element of Q_+ and determine the inverse of nonzero element $a \in Q_+$.

[Turn over

(d) Define an operation $*$ on the set of real numbers \mathbb{R} as $a * b = a + 2b$, $\forall a, b \in \mathbb{R}$. Then $(\mathbb{R}, *)$ is not a group because

- (i) \mathbb{R} is not closed w.r.t $*$
- (ii) \mathbb{R} is not associative w.r.t $*$
- (iii) Identity element does not exist.
- (iv) Inverse of each nonzero element does not exist.

(e) The value of i^i is

(i) $e^{-(4n+1)\frac{\pi}{2}}$ (ii) $e^{(4n-1)\frac{\pi}{2}}$

(iii) $e^{(4n+1)\frac{\pi}{2}}$ (iv) $e^{-(4n-1)\frac{\pi}{2}}$

(f) If z is a complex number, then $\sin^{-1}z$ is

(i) $-i \log \left(iz \pm \sqrt{1-z^2} \right)$

(ii) $i \log \left(iz \pm \sqrt{1+z^2} \right)$

$$(iii) -i \log \left(z \pm i\sqrt{1-z^2} \right)$$

$$(iv) i \log \left(z \pm i\sqrt{1+z^2} \right)$$

(g) If A is a square matrix, then $\text{adj } A' - (\text{adj } A)'$ is

(i) $2 |A|$

(ii) $2 |A| I$

(iii) null matrix

(iv) identity matrix

(h) Inverse of which of the following matrices exists? Given that $1, \omega, \omega^2$ are the cube roots of unity.

(i) $\begin{pmatrix} 1 & \omega \\ \omega & \omega^2 \end{pmatrix}$

(ii) $\begin{pmatrix} \omega^2 & 1 \\ 1 & \omega \end{pmatrix}$

(iii) $\begin{pmatrix} \omega & \omega^2 \\ \omega^2 & 1 \end{pmatrix}$

(iv) $\begin{pmatrix} \omega & \omega^2 \\ \omega & 1+\omega \end{pmatrix}$

(i) The rank of the matrix

$$\begin{pmatrix} 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \end{pmatrix} \text{ is}$$

(i) 0 (ii) 1

(iii) 2 (iv) 3

(j) If $2 + \sqrt{3}$ is a root of the equation $x^4 + 2x^3 - 16x^2 - 22x + 7 = 0$, then the other three roots are

(i) $2 - \sqrt{3}, 3 + \sqrt{2}, -3 + \sqrt{2}$

(ii) $-2 - \sqrt{3}, -3 + \sqrt{2}, -3 - \sqrt{2}$

(iii) $2 - \sqrt{3}, -3 + \sqrt{2}, -3 - \sqrt{2}$

(iv) $-2 + \sqrt{3}, 3 + \sqrt{2}, -3 - \sqrt{2}$

2. Give answers to the following questions :

$$2 \times 5 = 10$$

(a) If $f : A \rightarrow B$ and $g : B \rightarrow C$ are bijective mappings, then prove that $g \circ f$ is also a bijective mapping.

(b) In a group G , prove that $(ab)^{-1} = b^{-1} a^{-1}$,
 $\forall a, b \in G$.

(c) Show that every real symmetric matrix is Hermitian.

(d) Prove that

$$\pi = 2\sqrt{3} \left(1 - \frac{1}{3 \cdot 3} + \frac{1}{5 \cdot 3^2} - \frac{1}{7 \cdot 3^3} + \dots \right).$$

(e) If α, β, γ are the roots of the equation $x^3 + px^2 + qx + r = 0$, find the value of

$$\sum \frac{1}{\alpha^2 \beta^2}.$$

3. Answer any four parts :

5×4=20

(a) Define an equivalence relation on a nonempty set.

Show that the relation "congruence modulo m " is an equivalence relation on the set of integers.

1+4=5

(b) Let $f : A \rightarrow B$, $g : B \rightarrow C$, $h : C \rightarrow D$ be three mappings. Prove that

(i) $h \circ (g \circ f) = (h \circ g) \circ f$

(ii) $f \circ i = f$ and $j \circ f = f$ where $i : A \rightarrow A$ and $j : B \rightarrow B$ are identity mappings.

3+2=5

(c) If G is a group in which $(ab)^i = a^i b^i$ for three consecutive integers i and any a, b in G , then prove that G is abelian.

(d) (i) Find the values of $(1+i)^{\frac{1}{3}}$

(ii) Prove that

$$\sin^2 \theta \cos \theta = \theta^2 - \frac{5}{6} \theta^4 + \dots$$

$$+ (-1)^{n+1} \frac{3^{2n} - 1}{4(2n)!} \theta^{2n} + \dots \quad 2+3=5$$

(e) (i) Apply Descartes's rule of signs to ascertain the minimum number of complex roots of the equation $x^7 - 3x^3 + x + 1 = 0$.

(ii) If α, β, γ are the roots of the equation $x^3 + px^2 + qx + r = 0$, find the equation whose roots are $\alpha\beta + \beta\gamma, \beta\gamma + \gamma\alpha$ and $\gamma\alpha + \alpha\beta$. 3+2=5

(f) Define rank of a matrix. Find the rank of the matrix 1+4=5

$$\begin{pmatrix} 2 & 3 & -1 & -1 \\ 1 & -1 & -2 & -4 \\ 3 & 1 & 3 & -2 \\ 6 & 3 & 0 & -7 \end{pmatrix}$$

4. Answer any *one* part : 10

(a) (i) Prove that if G is a finite group and H is a subgroup of G , then $o(H)$ divides $o(G)$.

(ii) If G is a finite group, show that for any $a \in G$, $a^{o(G)} = e$ where e is the identity element of the group G .

(iii) Prove that every quotient group of a cyclic group is cyclic. $4+2+4=10$

(b) (i) Prove that centre of a group G is a subgroup of the group G .

(ii) For any integer a and a prime p , prove that $a^p \equiv a \pmod{p}$.

(iii) Prove that a finite group of prime order is abelian. $3+3+4=10$

5. Answer any *one* part : 10

(a) (i) If z is a complex number, show that

$$|z| \geq \frac{1}{\sqrt{2}} (|\operatorname{Re} z| + |\operatorname{Im} z|).$$

(ii) Express $\log(x + iy)$ in the form $A + iB$ where A and B are reals. Also find $\log(x + iy)$.

(iii) If $\cos^{-1}(\alpha + i\beta) = \theta + i\phi$, prove that $\alpha^2 \sec^2 \theta + \beta^2 \operatorname{cosec}^2 \theta = 1$

$$3+4+3=10$$

(b) (i) Two complex numbers z_1 and z_2 are such that $|z_1 + z_2| = |z_1 - z_2|$. Show that amp

z_1 and amp z_2 differ by $\frac{\pi}{2}$ or $\frac{3\pi}{2}$.

(ii) Expand $\tan x$ in ascending powers of x .

(iii) Deduce the Gregory series

$$\theta = \tan \theta - \frac{1}{3} \tan^3 \theta + \frac{1}{5} \tan^5 \theta - \dots \infty,$$

$$\text{where } -\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}.$$

$$3+3+4=10$$

6. Answer any two parts : $5 \times 2 = 10$

(a) Find the value of k and solve the equation $8x^3 - 12x^2 - kx + 3 = 0$ if the roots are in arithmetic progression. 5

(b) Solve the equation $x^3 - x^2 + 3x - 27 = 0$
if it have three distinct roots of equal moduli.

5

(c) Solve the equation $x^4 - 4x^3 - 4x^2 - 4x - 5 = 0$
given that two roots are connected by the
relation $2\alpha + \beta = 3$.

5

(d) Solve the equation $x^3 - 3x - 1 = 0$ by Cardon's
method.

5

7. Answer any *two* parts : 5×2=10

(a) Define a symmetric and a skew-symmetric
matrices.

Show that the matrix $B'AB$ is symmetric or
skew-symmetric according as A is symmetric
or skew-symmetric.

2+3=5

(b) If A is a square matrix of order n , prove that

$$|\text{adj}(\text{adj}A)| = |A|^{(n-1)^2}$$

5

(c) Prove that the necessary and sufficient
condition for a matrix A to possess an
inverse is that $|A| \neq 0$.

5

(d) For what values of λ the equations

$$x + y + z = 1$$

$$x + 2y + 4z = \lambda$$

$$x + 4y + 10z = \lambda^2$$

have a solution and solve them completely in each case.

5

Total No. of printed pages = 7

3 (Sem 1) MAT M2

2015

MATHEMATICS

(Major)

Paper : 1.2

(Calculus)

Full Marks = 80

Time - Three hours

The figures in the margin indicate full marks for the questions.

1. Answer the following : 1×10=10

(a) Write down the n th derivative of $\log_e(ax+b)$.

(b) If $u = \sin^{-1} \frac{x}{y} + \tan^{-1} \frac{y}{x}$, find $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y}$.

(c) Find $\frac{ds}{dx}$ for the curve $y^2 = 4ax$.

[Turn over