

2016

MATHEMATICS

( Major )

Paper : 5.1

( Real and Complex Analysis )

Full Marks : 60

Time : 3 hours

*The figures in the margin indicate full marks for the questions*

1. Answer the following questions : 1×7=7

- (a) State a sufficient condition for the continuity of a real-valued function of two variables.
- (b) Give an example of a real-valued function which is bounded but not Riemann integrable.
- (c) A real-valued function  $f$  is defined on  $[a, b]$  having a singular point in its domain. State whether  $f$  is Riemann integrable or not.

- (d) A function  $f(z) = u(x, y) + iv(x, y)$  is defined such that

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

State whether  $f$  is analytic or not.

- (e) Evaluate

$$\frac{1}{2\pi i} \oint_C \frac{\cos \pi z}{z^2 - 9} dz$$

where  $C$  is a closed rectangle with vertices at  $z = 2 \pm i, -2 \pm i$ .

- (f) State Cauchy's integral formula.

- (g) Define conformal mapping.

2. Answer the following questions :

2×4=8

- (a) Discuss the continuity of the following function at  $(0, 0)$  :

$$\begin{aligned} f(x, y) &= \frac{xy^3}{x^2 + y^6}, \quad (x, y) \neq (0, 0) \\ &= 0, \quad (x, y) = (0, 0) \end{aligned}$$

- (b) Show that the integral

$$\int_0^1 x^{m-1} e^{-x} dx$$

is convergent for  $m > 0$ .

- (c) Prove that if  $w = f(z) = u + iv$  is analytic in a region  $R$ , then

$$\frac{dw}{dz} = \frac{\partial w}{\partial x} = -i \frac{\partial w}{\partial y}$$

where  $u, v$  are functions of two variables  $x, y$ .

- (d) Find the fixed points of the transformation  $w = \frac{2z-5}{z+4}$ .

3. Answer any *three* parts : 5×3=15

- (a) If

$$u = \cos x, v = \sin x \cos y$$

$$w = \sin x \sin y \cos z$$

then show that

$$\frac{\partial(u, v, w)}{\partial(x, y, z)} = (-i)^3 \sin^3 x \sin^2 y \sin z$$

The symbols have their usual meanings.

- (b) Prove that if  $f$  is a bounded function on  $[a, b]$ , then to every  $\epsilon > 0$ , there corresponds  $\delta > 0$  such that

$$U(p, f) < \int_a^b f dx + \epsilon$$

The symbols have their usual meanings.

( 4 )

(c) Show that the integral

$$\int_0^{\pi/2} \log \sin x \, dx$$

is convergent. Hence evaluate it.

(d) Prove that if  $f(z)$  and  $g(z)$  are analytic at  $z_0$  and  $f(z_0) = g(z_0) = 0$  but  $g'(z_0) \neq 0$ , then

$$\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = \frac{f'(z_0)}{g'(z_0)}$$

(e) Prove that if  $f(z)$  is integrable along a curve  $C$  having finite length  $L$  and if there exists a positive number  $M$  such that  $|f(z)| \leq M$  on  $C$ , then

$$\left| \int_C f(z) \, dz \right| \leq ML$$

4. Answer either (a) or (b) :

(a) (i) If  $v$  is a function of two variables  $x$  and  $y$ , and  $x = r \cos \theta$ ,  $y = r \sin \theta$ , then prove that

$$\frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = \frac{\partial^2 v}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2 v}{\partial \theta^2} + \frac{1}{r} \frac{\partial v}{\partial r} \quad 5$$

- (ii) Find the shortest distance from the origin to the hyperbola

$$x^2 + 8xy + 7y^2 = 225, z = 0 \quad 5$$

- (b) (i) Verify the convergence of the integral

$$\int_0^{\infty} \frac{x \tan^{-1} x}{(1+x^4)^{1/3}} dx \quad 5$$

- (ii) Find the value of  $p$  such that

$$\int_1^{\infty} \frac{\sin x}{x^p} dx$$

converges absolutely. 5

5. Answer either (a) or (b) :

- (a) (i) Prove that if a function  $f$  is Riemann integrable on  $[a, b]$ , then  $f^2$  is also Riemann integrable on  $[a, b]$ . 5

- (ii) A function  $f$  is defined on  $[-1, 1]$  as follows :

$$f(x) = 1, x \neq 0$$

$$= 0, x = 0$$

Show that  $f$  is integrable on  $[-1, 1]$  and calculate its value. 5

( 6 )

- (b) (i) Show that the function  $f$  defined as follows

$$f(x) = \frac{1}{2^n}, \text{ when } \frac{1}{2^{n+1}} < x \leq \frac{1}{2^n} \quad (n = 0, 1, 2, \dots)$$
$$= 0, \quad x = 0$$

is integrable on  $[0, 1]$ . Also evaluate

$$\int_0^1 f \, dx \quad 5$$

- (ii) If  $f$  and  $g$  are both differentiable on  $[a, b]$  and if  $f'$ ,  $g'$  are both integrable on  $[a, b]$ , then show that

$$\int_a^b f(x) g'(x) \, dx = [f(x) g(x)]_a^b - \int_a^b g(x) f'(x) \, dx \quad 5$$

6. Answer either (a) or (b) :

- (a) (i) If

$$u_1(x, y) = \frac{\partial u}{\partial x}$$

$$\text{and } u_2(x, y) = \frac{\partial u}{\partial y}$$

then prove that

$$f'(z) = u_1(z, 0) - i u_2(z, 0) \quad 5$$

- (ii) Prove that  $\frac{d}{dz}(z^2 \bar{z})$  does not exist anywhere. 5

( 7 )

(b) (i) Evaluate

$$\oint_C \bar{z}^2 dz$$

around the circle  $|z-1|=1$ .

5

(ii) Find a bilinear transformation which maps  $z=0, -i, -1$  into  $w=i, 1, 0$  respectively.

5

\*\*\*