

Total number of printed pages -8

**63 (FY)SEM-2/MIN2/MATMIN1024**

**2025**

**MATHEMATICS**

**(MINOR)**

Paper : MATMIN1024

***(Integral Calculus and Differential Equations)***

*Full Marks : 70*

*Pass Marks : 28*

Time : Three hours

***The figures in the margin indicate full marks for the questions.***

1. Answer the following questions :  $1 \times 10 = 10$

(i)  $\int e^{-\log x} dx$  is equal to

(a)  $-e^{-\log x} + C$

(b)  $-xe^{-\log x} + C$

(c)  $\log |x| = C$

(d) None of the above

(ii)  $\int \frac{dx}{\sqrt{x^2 - a^2}}$  is

(a)  $\log \frac{x + \sqrt{x^2 - a^2}}{a} + C$

(b)  $\log \frac{x - \sqrt{x^2 - a^2}}{a} + C$

(c)  $\log \frac{x + \sqrt{x^2 - a^2}}{x} + C$

(d)  $\log \frac{x - \sqrt{x^2 - a^2}}{x} + C$

(iii)  $\int_0^{\pi/2} \sin^n x \, dx$  is equal to

(a)  $\int_0^{\pi/2} \sin^n x \, dx$

(b)  $\int_0^{\pi/2} \cos^n x \, dx$

(c)  $-\int_0^{\pi/2} \cos^n x \, dx$

(d) None of the above

(iv) The integrating factor for the linear differential equation of first order

$$\frac{dy}{dx} + P(x)y = Q(x) \text{ is}$$

(a)  $e^{\int P dx}$

(b)  $e^{-\int P dx}$

(c)  $e^{\int P/Q dx}$

(d)  $e^{\int Q dy}$

(v) The general solution of the differential

$$\text{equation } \frac{d^2y}{dx^2} + 4\frac{dy}{dx} + 4y = 0 \text{ is}$$

(a)  $y = C_1 e^{2x} + C_2 e^{-2x}$

(b)  $y = C_1 e^{-2x} + C_2 e^{-2x}$

(c)  $y = C_1 \cos 2x + C_2 \sin 2x$

(d)  $y = C_1 \cos 2x + C_2 x \cos 2x$

(vi) The particular integral of the differential

$$\text{equation } y'' - 5y' + 4y = 0 \text{ is}$$

(a) 0

(b)  $y = 4e^x + e^{4x}$

(c)  $y = C_1 e^x + C_2 e^{4x}$

(d)  $y = 2x$

(vii) If  $y = Ae^{Bx+C}$  is the solution of homogeneous partial differential equation, then the order of the differential equation is

- (a) 1
- (b) 2
- (c) 3
- (d) 4

(viii) Which of the following is the general form of Cauchy-Euler equation for a second order linear differential equation ?

- (a)  $ax^2y'' + bxy' + cy = 0$
- (b)  $ax^2y'' + bxy' + c = 0$
- (c)  $ax^2y'' + by' + c = 0$
- (d)  $ay'' + bxy + cy' = 0$

(ix) Which of the following will be Wronskian of two functions ?

- (a)  $W(f, g) = fg' - f'g$
- (b)  $W(f, g) = ff' - gg'$

(c)  $W(f, g) = fg' + f'g$

(d)  $W(f, g) = f'g' - fg$

(x) The order of partial differential equation  $r^2 + 6S - t = 0$  (symbols have their usual meaning) is

- (a) One
- (b) Two
- (c) Three
- (d) Four

2. Answer the following questions : **(any five)**  
2×5=10

(i) Evaluate  $\int_0^{n/2} \sin^7 x dx$

(ii) Evaluate  $\int_0^{\infty} \frac{dx}{1+x^2}$

(iii) Find the integrating factor of the differential equation

$$x^2y dx - (x^3 + y^3)dy = 0$$

(iv) Solve  $(2x^3 + 4y)dx + (4x + y - 1)dy = 0$

(v) Solve  $(D^2 - 8D + 16)y = 0$

(vi) Solve  $p^2 + p - 6 = 0$  where  $p = \frac{dy}{dx}$ .

3. Answer **any six** from the following questions : 5×6=30

(i) Evaluate  $\int \frac{dx}{\sqrt{7-6x-x^2}}$

(ii) If  $\phi(n) = \int_0^{\pi/4} \tan^n x dx$ , show that

$$\phi(n) + \phi(n-2) = \frac{1}{n-1}$$

(iii) Show that  $x(x-1)^{-1}$  is an integrating factor of the equation

$$x(x-1)\frac{dy}{dx} - y = x^2(x-1)^2 \quad \text{and hence solve it.}$$

(iv) Solve

$$(2xy + y - \tan y)dx +$$

$$(x^2 - x \tan^2 y + \sec^2 y)dy = 0$$

(v) Find the complete primitive of  $x^2(y - px) = p^2y$

(vi) Solve  $(D^2 - 2D + 5)y = e^{2x} \sin x$

(vii) Form the partial differential equation by eliminating the function  $f$  and  $F$  from  $y = f(x - at) + F(x + at)$

(viii) Show that  $e^{2x}$  and  $e^{3x}$  are linearly independent solution of

$$\frac{d^2y}{dx^2} - 5\frac{dy}{dx} + 6y = 0$$

(ix) Evaluate  $\int_0^{\infty} \frac{dx}{(x^2 + 1)^2}$

4. Answer the following questions : **(any two)**  
10×2=20

(a) Define Cauchy-Euler equation of homogeneous linear differential equation and also solve 2+8=10

$$x^3 \frac{d^3y}{dx^3} + 2x^2 \frac{d^2y}{dx^2} + 2y = 10 \left( x + \frac{1}{x} \right)$$

(b) (i) Solve by the method of variation of parameter  $\frac{d^2y}{dx^2} + y = \operatorname{cosec} x$  5

(ii) Solve  $\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + 10y = 0$

given that  $y(0) = 4, \frac{dy}{dx} = 1$

at  $x = 0$ . 5

(c) (i) Obtain the reduction formula for  $\int \cos^n x dx$  5

(ii) Evaluate  $\int \sec^3 x dx$  5

(d) Prove that the necessary and sufficient condition that a differential equation  $Mdx + Ndy = 0$  is exact is that

5+5=10

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

and hence solve

$$(x^3 + 3xy^2)dx + (y^3 + 3x^2y)dy = 0$$